

# Know Your Treatment and Prevention Options for COVID-19

There are more treatment options available now for COVID-19. While most people do not need treatment, treatments can prevent hospitalization and death for persons who are high risk. **Talk to your healthcare provider** if you are at risk of serious illness and interested in preventive therapy OR if you are at high risk and test positive for COVID-19.



Treatments are not a substitute for vaccination. Vaccination remains the safest, most effective way to protect you from COVID-19. Treatments must be started early, so don't delay testing. Here is information on the available COVID-19 treatments.

	Evusheld	Paxlovid	Remdesivir	Bebtelovimab	Molnupiravir
Who can get it?	<ul style="list-style-type: none"> <li>• Ages 12+</li> <li>• Persons <b>without</b> COVID-19 infection OR recent exposure; <b>AND who:</b></li> <li>• Are immunocompromised,</li> <li>• Have received treatment that weakens the immune system, <b>OR</b></li> <li>• Can't get vaccinated due to severe allergic reaction.</li> </ul>	<ul style="list-style-type: none"> <li>• Ages 12+</li> <li>• Persons with COVID-19 who are at high risk of serious illness</li> </ul>	<ul style="list-style-type: none"> <li>• Adults and children 28 days of age and older and over 3.0 kg (7lbs.)</li> <li>• Persons with COVID-19 who are not hospitalized but are at high risk of serious illness</li> </ul>	<ul style="list-style-type: none"> <li>• Ages 12+</li> <li>• Persons with COVID-19 who are at high risk of serious illness</li> </ul>	<ul style="list-style-type: none"> <li>• Ages 18+</li> <li>• Persons with COVID-19 who are at high risk of serious illness</li> </ul>
When do I need to get it?	<b>Before</b> infection to help prevent COVID-19	Within <b>5 days</b> from start of symptoms	Within <b>7 days</b> from start of symptoms	Within <b>7 days</b> from start of symptoms	Within <b>5 days</b> from start of symptoms
How is it given?	<b>Injection</b>	<b>Pills</b> taken orally for 5 days	Once daily <b>Intravenous Infusion (IV) for 3 days</b>	One-time <b>Intravenous Infusion (IV)</b>	<b>Pills</b> taken orally for 5 days
Where can it be given?	<b>Healthcare Facility/ Infusion Center</b>	<b>Home</b>	<b>Healthcare Facility/ Infusion Center</b>	<b>Healthcare Facility/ Infusion Center</b>	<b>Home</b>

For more information, visit: <https://covid-19.acgov.org/treatment>

