Know Your Treatment and Prevention Options for COVID-19

There are more treatment options available now for COVID-19. While most people do not need treatment, treatments can prevent hospitalization and death for persons who are high risk. **Talk to your healthcare provider** if you are at risk of serious illness and interested in preventive therapy OR if you are at high risk and test positive for COVID-19.

Treatments are not a substitute for vaccination. Vaccination remains the safest, most effective way to protect you from COVID-19. Treatments must be started early, so don't delay testing. Here is information on the available COVID-19 treatments.

	Evusheld	Paxlovid	Molnupiravir	Bebtelovimab	Remdesivir
Who is eligible?	 Ages 12+ Persons without COVID-19 infection OR recent exposure; AND who: Are immunocompromised, Have received treatment that lowers the immune system, OR Can't get vaccinated due to severe allergic reaction. 	Ages 12+ Persons with COVID-19 who are at high risk of serious illness	Ages 18+ Persons with COVID-19 who are at high risk of serious illness	Ages 12+ Persons with COVID-19 who are at high risk of serious illness	 Adults and children over 3.5 kg (8lbs.) Persons with COVID-19 who are not hospitalized but are at high risk of serious illness
When must it be started?	Before infection to help prevent COVID-19	Within 5 days from start of symptoms	Within 5 days from start of symptoms	Within 7 days from start of symptoms	Within 7 days from start of symptoms
How is it given?	Injection	Pills taken orally for 5 days	Pills taken orally for 5 days	One-time Intravenous Infusion (IV)	Once daily Intravenous Infusion (IV) for 3 days
Where can it be given?	Healthcare Facility/ Infusion Center	Home	Home	Healthcare Facility/ Infusion Center	Healthcare Facility/ Infusion Center

For more information, visit: https://www.fda.gov/consumers/consumer-updates/know-your-treatment-options-covid-19

